

## Drug Facts

<b>Active ingredient (in each tablet)</b>	<b>Purpose</b>
Triprolidine HCl 2.5 mg .....	Antihistamine

### Uses

temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes

### Warnings

**Ask a doctor before use if you have**

- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor before use if you are taking sedatives or tranquilizers.**

**When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages while taking this product
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if new symptoms occur.** ▶

## Drug Facts (continued)

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- **do not take more than directed**
- **adults and children 12 years of age and over:** 1 tablet (2.5 mg) every 4 to 6 hours; not to exceed 4 tablets (10 mg) in 24 hours or as directed by a doctor
- **children under 12 years of age:** do not use

### Other information

- store at controlled room temperature 20-25°C (68-77°F)

**Inactive ingredients** hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, titanium dioxide

**Questions or comments?** 1-844-356-6273

**KEEP CARTON FOR COMPLETE INFORMATION.  
Do not use if blister is broken or damaged.**

\*Dosage is 1 tablet every 4 to 6 hours